arXiv:1607.04873v1 [math.AC] 17 Jul 2016

UNIFORM DETERMINANTAL REPRESENTATIONS

ADA BORALEVI*, JASPER VAN DOORNMALEN†, JAN DRAISMA†‡,
MICHIEL E. HOCHSTENBACH†, AND BOR PLESTENJAK§

Abstract. The problem of expressing a specific polynomial as the determinant of a square matrix of affine-linear forms arises from algebraic geometry, optimisation, complexity theory, and scientific computing. Motivated by recent developments in this last area, we introduce the notion of a *uniform determinantal representation*, not of a single polynomial but rather of all polynomials in a given number of variables and of a given maximal degree. We derive a lower bound on the size of the matrix, and present a construction achieving that lower bound up to a constant factor as the number of variables is fixed and the degree grows. This construction marks an improvement upon a recent construction due to Plestenjak–Hochstenbach, and we investigate the performance of new representations in their root-finding technique for bivariate systems. Furthermore, we relate uniform determinantal representations to vector spaces of singular matrices, and we conclude with a number of future research directions.

Key words. Determinantal representation, system of polynomial equations, multiparameter matrix eigenvalue problem, space of singular matrices.

AMS subject classifications. 13P15, 65H04, 65F15, 65F50.

1. Introduction and results.

Consider an n-variate polynomial of degree at most d:

$$p = \sum_{|\alpha| \leq d} c_\alpha x^\alpha$$

where $x := (x_1, \ldots, x_n)$, $\alpha \in \mathbb{Z}^n_{\geq 0}$, $|\alpha| := \sum_i \alpha_i$, $x^\alpha := \prod_i x_i^{\alpha_i}$, and where each coefficient c_α is taken from a ground field K. A *determinantal representation* of p is an $N \times N$-matrix M of the form

$$M = A_0 + \sum_{i=1}^{n} x_i A_i,$$

where each $A_i \in K^{N \times N}$, with $\det(M) = p$. We call N the *size* of the determinantal representation. Clearly, since the entries of M are affine-linear forms in $x_1, \ldots, x_n$, N must be at least the degree of p.

Determinantal representations of polynomials play a fundamental role in several mathematical areas: from *algebraic geometry* it is known that each plane curve ($n = 2$) of degree d over an algebraically closed field K admits a determinantal representation of size d [8, 10]. Over non-algebraically closed fields, and especially when restricting to symmetric determinantal representations, the situation is much more subtle [20]. For larger n, only certain hypersurfaces have a determinantal representation of size equal to their degree [2, 8]. In *optimisation*, and

*Scuola Internazionale Superiore di Studi Avanzati, via Bonomea 265, 34136 Trieste, Italy, ada.boralevi@sissa.it. This author is member of GNSAGA and was partially supported by PRIN 2010-2011 project "Geometria delle varietà algebriche"and Università di Trieste–FRA 2011 project "Geometria e topologia delle varietà".

†Department of Mathematics and Computer Science, TU Eindhoven, PO Box 513, 5600 MB, The Netherlands, m.j.v.doornmalen@student.tue.nl, j.draisma@tue.nl, m.e.hochstenbach@tue.nl. The first, third, and fourth author have been (partially) supported by NWO Vidi research grants.

‡Faculteit Exacte Wetenschappen, Afdeling Wiskunde, Vrije Universiteit, De Boelelaan 1081a, 1081 HV Amsterdam, The Netherlands.

§IMFM and Department of Mathematics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia, bor.plestenjak@fmf.uni-lj.si. This author was partially supported by the ARRS Grant P1-0294.

notably in the theory of *hyperbolic polynomials* [39], one is particularly interested in the case where $K = \mathbb{R}$, A_0 is symmetric positive definite, and the A_i are symmetric. In this case, the restriction of p to any line through 0 has only real roots. For $n = 2$ the converse also holds [16, 22]; for counterexamples to this converse holding for higher n, see [7]. In *complexity theory* a central role is played by Valiant's conjecture that the permanent of an $m \times m$-matrix does not admit a determinantal representation of size polynomial in m [37]. Via the *geometric complexity theory* programme [27] this leads to the study of polynomials in the boundary of the orbit of the $N \times N$-determinant under the action of the group $\mathrm{GL}_{N^2}(K)$ permuting matrix entries. Recent developments in this field include the study of this boundary for $N = 3$ [18] and the exciting negative result in [6] that Valiant's conjecture can *not* be proved using occurrence obstructions proposed earlier in [28].

Our motivation comes from *scientific computing*, where determinantal representations of polynomials have recently been proposed for efficiently solving systems of equations [31]. For this application, it is crucial to have determinantal representations not of a *single* polynomial p, but rather of all n-variate polynomials of degree at most d. Moreover, the representation should be easily computable from the coefficients of p. Specifically, in [31] determinantal representations are constructed for the bivariate case ($n = 2$) in which the entries of the matrices $A_0, \ldots, A_n$ themselves *depend affine-linearly* on the coefficients c_α. This is what we call a *uniform determinantal representation* of the generic polynomial p of degree d in n variables; see Section 2 for a precise definition.

Example 1.1 (The binary quadric). The identity

$$c_{00} + c_{10}x + c_{01}y + c_{20}x^2 + c_{11}xy + c_{02}y^2 = \det \begin{bmatrix} -x & 1 & 0 \\ -y & 0 & 1 \\ c_{00} & c_{10} + c_{20}x + c_{11}y & c_{01} + c_{02}y \end{bmatrix}$$

exhibits the matrix on the right as a uniform determinantal representation of the generic bivariate quadric. ♣

In applications, the matrix M is used as input to algorithms in numerical linear algebra that scale unfavourably with N, such as a complexity of $O(N^6)$. Consequently, we are led to consider the following fundamental question.

Question 1.2. *What is the minimal size $N^*(n, d)$ of any uniform determinantal representation of the generic polynomial of degree d in n variables?*

A construction from [31] shows that for fixed $n = 2$ and $d \to \infty$ we have $N^*(2, d) \leq \frac{1}{4}\, d^2 + O(d)$; this construction is reviewed in Section 4. We improve the construction from [31] by giving a particularly elegant uniform determinantal representations of bivariate polynomials of size $2d + 1$ in Example 5.1, and of size $2d - 1$ in Example 5.3. In view of the obvious lower bound of d this is clearly sharp up to a constant factor for $d \to \infty$, although we do not know where in the interval $[d, 2d - 1]$ the true answer lies. We show in Section 8 how to use these small determinantal representations of bivariate polynomials for solving systems of equations. Before that, we focus on the asymptotic behaviour of $N^*(n, d)$ for fixed n and $d \to \infty$. In this setting, we derive the following result.

Theorem 1.3. *For fixed $n \in \mathbb{Z}_{\geq 2}$ there exist positive constants C_1, C_2 (depending on n) such that for each $d \in \mathbb{Z}_{\geq 0}$ the smallest size $N^*(n, d)$ of a uniform determinantal representation of the generic polynomial of degree d in n variables satisfies $C_1 d^{n/2} \leq N^*(n, d) \leq C_2 d^{n/2}$.*

We will also compare our results with previous constructions, most notably with those by Quarez [33, Thm. 4.4], who proves the existence of a symmetric representation of size $\binom{n+\lfloor \frac{d}{2} \rfloor}{n}$. For fixed n and $d \to \infty$, [33] therefore has the asymptotic rate $\sim d^n$, meaning that the results of

this paper represent a clear improvement. For fixed d and $n \to \infty$, [33] leads to the asymptotic behavior $\sim n^{\lfloor d/2 \rfloor}$, which is similar to our bounds; we will discuss more details in Section 9.

In Section 2 we formalise the notion of uniform determinantal representations, study their symmetries, and derive some simple properties. In particular, we relate uniform determinantal representations to spaces of singular $N \times N$-matrices. In Section 3 we briefly review some of the existing literature on these singular spaces, and we prove that for $N > 4$ there are infinitely many equivalence classes of such objects; this poses an obstruction to a "brute-force" approach towards finding lower bounds on $N^*(n, d)$. In Section 4 we present a first construction, of which however the size is of the order of d^n, rather than $d^{n/2}$, for $d \to \infty$. In Section 5 we give a more efficient construction and prove Theorem 1.3. In Section 6 we give upper bounds on $N^*(n, d)$ for small n and d and determine $N^*(2, 2)$ and $N^*(3, 2)$ exactly. We extend representations from scalar to matrix polynomials in Section 7. In Section 8 we give some numerical results that show that for $n = 2$ and small d we get a competitive method for computing zeros of polynomials systems. Finally, in Section 9 we summarise our main conclusions and collect some questions that arise naturally from our work.

2. Problem formulation and symmetries. In this section we give a formal definition of uniform determinantal representations, and introduce a group that acts on such representations. We also show that a uniform determinantal representation gives rise to a vector space consisting entirely of singular matrices; such spaces are the topic of next section.

Let K be a field and fix $d, n \in \mathbb{Z}_{\geq 0}$. Let F_d denote the polynomials of degree at most d in the polynomial ring $K[x_1, \ldots, x_n]$. Furthermore, let $p_{n,d}$ be the generic polynomial of that degree, i.e.,

$$p_{n,d} = \sum_{|\alpha| \leq d} c_\alpha x^\alpha, \tag{2.1}$$

where $x := (x_1, \ldots, x_n)$, $\alpha \in \mathbb{Z}^n_{\geq 0}$, $|\alpha| := \sum_i \alpha_i$, $x^\alpha := \prod_i x_i^{\alpha_i}$, and where we consider c_α as a variable for each α.

DEFINITION 2.1. *For $n, d \in \mathbb{Z}_{\geq 0}$, a* uniform determinantal representation *of $p_{n,d}$ is an $N \times N$-matrix M with entries from $K[(x_1, \ldots, x_n), (c_\alpha)_{|\alpha| \leq d}]$, of degree at most 1 in each of these two sets of variables, such that $\det(M) = p_{n,d}$. The number N is called the* size *of the determinantal representation.*

To be explicit, we require each entry of M to be a K-linear combination of the monomials $1, x_i, c_\alpha, c_\alpha x_i$, $(i = 1, \ldots, n, |\alpha| \leq d)$. This means that we can decompose M as $M_0 + M_1$, where M_0 contains all terms in M that do not contain any c_α, and where M_1 contains all terms in M that do. We will use the notation $M = M_0 + M_1$ throughout the paper. When n and d are fixed in the context, we will also speak of a uniform determinantal representation without reference to $p_{n,d}$. Our ultimate aim is to determine the following quantity.

DEFINITION 2.2. *For $n, d \in \mathbb{Z}_{\geq 0}$, $N^*(n, d) \in \mathbb{Z}_{>0}$ is the minimum among all sizes of uniform determinantal representations of $p_{n,d}$.*

This minimal size could potentially depend on the ground field K, but the bounds that we will prove do not. Note that in the definition of $N^*(n, d)$ we do not allow terms in M of degree strictly larger than one in the c_α. Relaxing this condition to polynomial dependence on the c_α might affect the exact value of N^*, but it will not affect our bounds—see Remark 2.7.

Given a uniform determinantal representation M of size N, and given matrices g, h in $\mathrm{SL}_N(K)$, the group of determinant-one matrices with entries in K, the matrix gMh^{-1} is another uniform determinantal representation of $p_{n,d}$. In this manner, the group $\mathrm{SL}_N(K) \times \mathrm{SL}_N(K)$ acts on the set of uniform determinantal representations of $p_{n,d}$. Moreover, there exist further symmetries, arising from affine transformations of the n-space. Recall that these transformations

form the group $\mathrm{AGL}_n(K) = \mathrm{GL}_n(K) \ltimes K^n$ generated by invertible linear transformations and translations.

LEMMA 2.3. *The group* $\mathrm{AGL}_n(K)$ *acts on uniform determinantal representations of* $p_{n,d}$.

The statement of this lemma is empty without making the action explicit, as we do in the proof.

Proof. Let $g \in \mathrm{AGL}_n(K)$ be an affine transformation of K^n, and expand

$$p_{n,d}(g^{-1}x, c) = \sum_{|\alpha| \leq d} c'_\alpha x^\alpha,$$

where the c'_α are linear combinations of the c_α. More precisely, the vector c' can be written as $\rho(g)c$, where ρ is the representation of $\mathrm{AGL}_n(K)$ on polynomials of degree at most d regarded as a matrix representation relative to the monomial basis.

Now let $M = M(x, c)$ be a uniform determinantal representation of $p_{n,d}$. Then

$$\det(M(g^{-1}x, \rho(g)^{-1}c)) = p_{n,d}(g^{-1}x, \rho(g)^{-1}c) = p_{n,d}(x, c),$$

i.e., $M(g^{-1}x, \rho(g)^{-1}c)$ is another uniform determinantal representation of $p_{n,d}$. The action of g is given by $M \mapsto M(g^{-1}x, \rho(g)^{-1}c)$. □

EXAMPLE 2.4 (The binary quadric revisited). Take $n = d = 2$ and the affine transformation $g(x, y) := (y, x + 1)$ with inverse $g^{-1}(x, y) = (y - 1, x)$. We have

$$p_{2,2}(g^{-1}(x, y)) = (c_{00} - c_{10} + c_{20}) + (c_{01} - c_{11})x + (c_{10} - 2c_{20})y + c_{02}x^2 + c_{11}xy + c_{20}y^2.$$

We find

$$c' = \begin{bmatrix} c'_{00} \\ c'_{10} \\ c'_{01} \\ c'_{20} \\ c'_{11} \\ c'_{02} \end{bmatrix} = \begin{bmatrix} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} c_{00} \\ c_{10} \\ c_{01} \\ c_{20} \\ c_{11} \\ c_{02} \end{bmatrix} = \rho(g)c; \ \rho(g)^{-1} = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}.$$

If we make the substitutions

$$\begin{array}{llll} c_{00} \mapsto c_{00} + c_{01} + c_{02} & c_{10} \mapsto c_{01} + 2c_{02} & c_{01} \mapsto c_{10} + c_{11} & x \mapsto y - 1 \\ c_{20} \mapsto c_{02} & c_{11} \mapsto c_{11} & c_{02} \mapsto c_{20} & y \mapsto x \end{array}$$

in the uniform determinantal representation of Example 1.1, then we arrive at the matrix

$$\begin{bmatrix} 1 - y & 1 & 0 \\ -x & 0 & 1 \\ c_{00} + c_{01} + c_{02} & c_{01} + c_{02} + c_{02}y + c_{11}x & c_{10} + c_{11} + c_{20}x \end{bmatrix}$$

whose determinant also equals $p_{2,2}$. ♣

The action of the affine group will be used in Section 6 to determine the exact value of $N^*(n, 2)$ for $n = 2$ and 3. We now turn our attention to the component M_0 of a uniform determinantal representation M.

LEMMA 2.5. *For any uniform determinantal representation* $M = M_0 + M_1$ *of size* N*, the determinant of* M_0 *is the zero polynomial in* $K[x_1, \ldots, x_n]$*. Moreover, at every point* $\bar{x} \in K^n$*, the rank of the specialisation* $M_0(\bar{x}) \in K^{N \times N}$ *is exactly* $N - 1$.

Proof. The first statement follows from the fact that $\det(M_0)$ is the part of the polynomial $\det(M)$ which is homogeneous of degree zero in the c_α; hence zero.

By specialising the vector x of variables to a point $\bar{x} \in K^n$, the rank of M_0 can only drop, so the rank of $M_0(\bar{x})$ is at most $N-1$. However, if it were at most $N-2$, then after column operations on M by means of determinant-one matrices with entries in K we may assume that $M_0(\bar{x})$ has its last two columns equal to 0. This means that all entries of $M(\bar{x}) = M_0(\bar{x}) + M_1(\bar{x})$ in these columns are linear in the c_α. This in turn implies that any term in the polynomial $\det M(\bar{x})$ is at least quadratic in the c_α. But on the other hand $\det M(\bar{x})$ equals $p_{n,d}(\bar{x})$, which is a non-zero *linear* polynomial in the c_α (nonzero since not every polynomial of degree at most d vanishes at $\bar{x}$). This contradiction implies that the rank of $M_0(\bar{x})$ is $N-1$. □

LEMMA 2.6. *If $M = M_0 + M_1$ is a uniform determinantal representation of size N, then $V \subseteq F_{N-1}$ spanned by the $(N-1)\times(N-1)$-subdeterminants of M_0 satisfies $F_1 \cdot V \supseteq F_d$.*

Here, as in the rest of this paper, by the product of two spaces of polynomials we mean the K-linear span of all the products.

Proof. Let D_{ij} be the determinant of the submatrix of M_0 obtained by deleting the ith row and the jth column. On the one hand, $\det(M) = p_{n,d}$ is linear in the c_α by assumption, and on the other hand, by expanding $\det(M)$ we see that the part that is homogeneous of degree one in the c_α is

$$\sum_{i,j} (-1)^{i+j} (M_1)_{ij} D_{ij};$$

this therefore equals $p_{n,d}$. Hence any element q of F_d is obtained from the expression above by specialising the variables c_α to the coefficients of q. Since each $(M_1)_{ij}$ is then specialised to an element of F_1, we find $q \in F_1 \cdot V$. □

REMARK 2.7. Note that the proof above still applies if we allow determinantal representations of the form $M_0 + M_1 + \cdots + M_e$ where M_r is homogeneous of degree r in the c_α and affine-linear in the x_i. Since our upper bound in Section 5 builds directly on this lemma, the upper bound holds in this more general setting, as well.

3. Spaces of singular matrices. Let $M = M_0 + M_1$ be a uniform determinantal representation of $p_{n,d}$. Writing $M_0 = B_0 + \sum_{i=1}^n x_i B_i$, Lemma 2.5 implies that the linear span $\langle B_0, \ldots, B_n \rangle_K \subseteq K^{N\times N}$ consists entirely of singular matrices (and indeed that this remains true when extending scalars from K to an extension field). There is an extensive literature on such *singular matrix spaces*; see, e.g., [11, 13] and the references therein. The easiest examples are the following.

DEFINITION 3.1. *A subspace $\mathcal{A} \subseteq K^{N\times N}$ is called a* compression space *if there exists a subspace $U \subseteq K^N$ with $\dim(\langle u^T A \mid A \in \mathcal{A}, u \in U\rangle_K) < \dim U$. We call the space U a* witness *for the singularity of $\mathcal{A}$.*

Given any two subspaces $U, V \subseteq K^N$ with $\dim V = -1 + \dim U$, the space of all matrices which map U into V (acting on row vectors) is a compression space with witness U. It is easy to see that these spaces are inclusion-wise maximal among all singular spaces.

If $\mathcal{A}$ is a singular matrix space, then so is $g\mathcal{A}h^{-1}$ for any pair $(g,h) \in \mathrm{GL}_N(K) \times \mathrm{GL}_N(K)$. We call the latter space *conjugate* to the former.

EXAMPLE 3.2. For $N = 2$, every singular matrix space is a compression space, hence conjugate to a subspace of one of the two spaces

$$\left\{ \begin{bmatrix} * & * \\ 0 & 0 \end{bmatrix} \right\}, \left\{ \begin{bmatrix} 0 & * \\ 0 & * \end{bmatrix} \right\},$$

where the $*$s indicate entries that can be filled arbitrarily. A witness for the first space is the span $\langle e_2 \rangle$ of the second standard basis vector, and a witness for the second space is K^2.

For $N = 3$, there are four conjugacy classes of inclusion-maximal singular matrix spaces, represented by the three maximal compression spaces

$$\left\{\begin{bmatrix} * & * & * \\ * & * & * \\ 0 & 0 & 0 \end{bmatrix}\right\}, \left\{\begin{bmatrix} * & * & * \\ 0 & 0 & * \\ 0 & 0 & * \end{bmatrix}\right\}, \left\{\begin{bmatrix} 0 & * & * \\ 0 & * & * \\ 0 & * & * \end{bmatrix}\right\},$$

and the space of skew-symmetric 3×3-matrices [12]; the latter is not a compression space. For $N = 4$, there are still finitely many (namely, 10) conjugacy classes of inclusion-maximal singular matrix spaces [12, 13], but this is not true for $N \geq 5$, as Theorem 3.4 below shows. This theorem is presumably folklore; we include a proof since we have not been able to find a literature reference for it. ♣

PROPOSITION 3.3. *Assume that K is algebraically closed. For any m and $N \in \mathbb{Z}_{\geq 0}$ the locus X_m in the Grassmannian $\mathrm{Gr}(m, K^{N\times N})$ of m-dimensional subspaces of $K^{N\times N}$ consisting of all* singular *subspaces is closed in the Zariski topology. Moreover, the locus U_m in X_m consisting of all* inclusion-wise maximal *singular subspaces is open inside X_m.*

Proof. The first statement is standard. For the second statement, consider the incidence variety

$$Z := \{(\mathcal{A}, \mathcal{A}') \in X_m \times X_{m+1} \mid \mathcal{A} \subseteq \mathcal{A}'\} \subseteq X_m \times X_{m+1},$$

which is a closed subvariety of $X_m \times X_{m+1}$. The projection of Z into X_m is the complement of U_m, and it is closed because X_{m+1} is a projective variety. □

THEOREM 3.4. *Assume that K is infinite and of characteristic unequal to two. For $N \geq 5$ there are infinitely many conjugacy classes of inclusion-wise maximal singular $N \times N$-matrix spaces.*

Proof. Take $N \geq 5$. For sufficiently general skew-symmetric matrices $A_1, \ldots, A_N \in K^{N\times N}$ set $A := (A_1, \ldots, A_N)$ and define the space

$$\mathcal{B}_A := \{(A_1x| \cdots |A_Nx) \mid x \in K^N\} \subseteq K^{N\times N}.$$

Each matrix in this space is singular, since for $x \neq 0$ we have

$$x^T(A_1x| \cdots |A_Nx) = (x^TA_1x, \ldots, x^TA_Nx) = 0.$$

In [13] it is proved that, for a specific choice of the tuple A, the space $\mathcal{B}_A$ is maximal among the singular subspaces of $K^{N\times N}$. By Proposition 3.3, $\mathcal{B}_A$ is maximal for sufficiently general A, as well (note that we may first extend K to its algebraic closure to apply the proposition). In the notation of that proposition, we have a rational map

$$\varphi : S^N \dashrightarrow U_N, \quad A \mapsto \mathcal{B}_A,$$

where $S \subseteq K^{N\times N}$ denotes the subspace of skew-symmetric matrices; the dashed arrow indicates that the map is defined only in an open dense subset of S^N. For any nonzero scalar t, $\varphi(tA) = \varphi(A)$. We claim that, in fact, the general fibre of φ is indeed one-dimensional. As the fibre dimension is semicontinuous, it suffices to verify this at a particular point where φ is defined. We take $A_i = E_{i,i+1} - E_{i+1,i}$ for $i = 1, \ldots, N-1$ and A_N general; here E_{ij} is the matrix with zeros everywhere except for a 1 at position (i, j). Let $B \in S^N$; if $\varphi(A) = \varphi(B)$, then there exists an invertible matrix $g \in \mathrm{GL}_N(K)$ such that

$$(A_1gx| \cdots |A_Ngx) = (B_1x| \cdots |B_Nx)$$

for all x, so that $A_i g = B_i$. Using skew-symmetry of A_i and B_i, we find that $A_i g = g^T A_i$. Substituting our choice of A_i for $i \in \{1, \ldots, N-1\}$ yields $g_{i,j} = g_{j,i} = 0$ for all j with $|i-j| > 1$, $g_{i,i+1} = -g_{i,i+1}$, so $g_{i,i+1} = 0$ since $\operatorname{char} K \neq 2$, and $g_{i,i} = g_{i+1,i+1}$. Hence g is a scalar multiple of the identity. It follows that the fibre of φ through A is one-dimensional as claimed.

Since $\dim S = \binom{N}{2}$, we have thus constructed an $(N\binom{N}{2} - 1)$-dimensional family inside U_N. Given any point $\mathcal{A}$ in U_N, its orbit under $\mathrm{GL}_N(K) \times \mathrm{GL}_N(K)$ has dimension at most $2(N^2-1)$ (scalars act trivially). Now for $N = 5$ we have

$$N\binom{N}{2} - 1 = 5 \cdot 10 - 1 = 49 \quad \text{and} \quad 2(N^2-1) = 48,$$

so that we have found (at least) a one-parameter family of conjugacy classes of singular spaces. For $N > 5$ the difference between $N\binom{N}{2} - 1$ and $2(N^2-1)$ is even larger. □

For large N it seems impossible to classify maximal singular matrix spaces. The construction above already gives an infinite number of conjugacy classes, but there are many other sources of examples. For instance, for infinitely many N there exists a maximal singular matrix space in $K^{N\times N}$ of constant dimension 8, at least if we assume that K has characteristic 0 [11]. On the other hand, if the singular matrix space $\mathcal{A}$ has dimension at least $N^2 - N$, then it is a compression space with either a one-dimensional witness or all of K^N as witness [9] (and hence of dimension exactly $N^2 - N$). A sharpening of this result is proved in [13] (see also [35]).

It should be noted that in many cases not even the dimension of such singular matrix spaces is known, for fixed values of the size and rank of the matrices. There is a considerable body of work devoted to giving lower and upper bounds for such dimensions, both in the case of bounded and constant rank, but these bounds are rarely sharp, see, among many other references, [14, 19, 36, 40] and the more recent works on skew-symmetric matrices of constant rank [4, 23].

Hence the fact that M_0 represents a singular matrix space of dimension (at most) $n+1$ does not much narrow down our search for good uniform determinantal representations, except in small cases discussed in Section 6. However, for our constructions in Sections 4 and 5 we will only use compression spaces where the witness has dimension 1 or about $\frac{1}{2}N$, respectively; and our lower bounds on $N^*(n,d)$ are independent of the literature on singular matrix spaces.

4. A first construction. In this section we restrict our attention to determinantal representations $M = M_0 + M_1$ where M_0 represents a compression space with a one-dimensional witness (or, dually by transposition, with a full-dimensional witness). Under this assumption we will prove quite tight bounds on the minimal size of a uniform matrix representation. The following fundamental notion will be used throughout below.

Definition 4.1. *We say that a subspace $V \subseteq K[x_1, \ldots, x_n]$ is* connected to 1 *if it is nonzero and its intersections $V_e := V \cap F_e$ satisfy $F_1 \cdot V_e \supseteq V_{e+1}$ for each $e \geq 0$.*

Note that this implies that $V_0 = \langle 1 \rangle$. We borrow the terminology from the theory of border bases [21], where a set S of monomials is called connected to 1 if $1 \in S$ and each nonconstant monomial in S can be divided by some variable to obtain another monomial in S. The linear span of S is then connected to 1 in our sense. Translating monomials to their exponent vectors, we will call a subset S of $\mathbb{Z}^n_{\geq 0}$ *connected to* 0 if it contains 0 and for each $\alpha \in S \setminus \{0\}$ there exists an i such that $\alpha - e_i \in S$, where e_i is the i-th standard basis vector.

Let V be a finite-dimensional subspace of $K[x_1, \ldots, x_n]$ connected to 1. Choose a K-basis $f_1, \ldots, f_m$ of V whose total degrees increase weakly. For each $i = 2, \ldots, m$ write

$$f_i = \sum_{j<i} \ell_{ij} f_j$$

for suitable elements $\ell_{ij} \in F_1$. Let M_V be the $(m-1) \times m$-matrix whose ith row equals

$$(-\ell_{i1}, -\ell_{i2}, \ldots, -\ell_{i,i-1}, 1, 0, \ldots, 0).$$

Note that M_V depends on the choice of basis, but we suppress this dependence in the notation, since the property of M_V in the next lemma does not depend on the choice of basis.

LEMMA 4.2. *The K-linear subspace of $K[x_1, \ldots, x_n]$ spanned by the $(m-1) \times (m-1)$-subdeterminants of M_V equals V.*

Proof. By construction, M_V has rank $m-1$ over the field $K(x_1, \ldots, x_n)$ and satisfies $M_V \cdot (f_1, \ldots, f_m)^T = 0$. By (a version of) Cramer's rule, the kernel of M_V is also spanned by $(D_1, -D_2, \ldots, (-1)^{m-1} D_m)$ where D_j is the determinant of the submatrix of M_V obtained by removing the jth column. So these two vectors differ by a factor in $K(x_1, \ldots, x_n)$. Since $D_1 = 1 = f_1$ we find that they are, in fact, equal. Hence $\langle D_1, \ldots, D_m \rangle = V$ as claimed. □

We can now formulate our first general construction. This generalises a construction from [31] to the multivariate case.

PROPOSITION 4.3. *Let V be an m-dimensional space connected to 1 and suppose that $F_1 \cdot V \supseteq F_d$. Then there exists a uniform determinantal representation of size m for the generic polynomial of degree d in n variables.*

Proof. Let $D_1, \ldots, D_m \in K[x_1, \ldots, x_n]$ be the $(m-1) \times (m-1)$-subdeterminants of M_V. By $F_1 \cdot V = F_d$ and Lemma 4.2 we can find, for each monomial x^α of degree at most d, affine-linear forms $l_{\alpha 1}, \ldots, l_{\alpha m} \in F_1$ such that $x^\alpha = \sum_j (-1)^{j-1} l_{\alpha j} D_j$. Define

$$s := \sum_{|\alpha| \leq d} c_\alpha (l_{\alpha 1}, \ldots, l_{\alpha m}),$$

a row vector of bi-affine linear forms in the x_i and the c_α. Then, by Laplace expansion along the first row, we find that the determinant of

$$\begin{bmatrix} s \\ M_V \end{bmatrix}$$

is the generic polynomial of degree d in $x_1, \ldots, x_n$. □

EXAMPLE 4.4. For $n = 2$ the following picture gives a space V, connected to 1 and spanned by the monomials marked with black vertices, such that $F_1 \cdot V = F_6$:



This is the construction of [31], which shows that there exists a uniform determinantal representation of the generic bivariate polynomial of degree d of size $\frac{1}{4} d^2 + O(d)$ as $d \to \infty$. ♣

The bivariate case generalises as follows.

THEOREM 4.5. *For fixed n, there exists a determinantal representation $M = M_0 + M_1$ of the generic n-variate polynomial of degree d of size $\frac{1}{n \cdot n!} d^n + O(d^{n-1})$ such that, moreover, the singular matrix space represented by M_0 is a compression space with a one-dimensional witness. Moreover, under this latter additional condition on M_0, the bound is sharp.*

Proof. Note that $\dim F_d = \binom{n+d}{n} = \frac{1}{n!} d^n + O(d^{n-1})$. Hence by Proposition 4.3 it suffices to show the existence of a subspace $V \subseteq F_d$ connected to 1 and such that $F_1 \cdot V = F_d$, where

$\dim V = \frac{1}{n} \dim F_d + O(d^{n-1})$. We will, in fact, show that V can be chosen to be spanned by monomials.

First, recall that there exists a lattice Λ in $\mathbb{Z}^{n-1}$ such that $\mathbb{Z}^{n-1}$ is the disjoint union of Λ and its cosets $e_i + \Lambda$ for $i = 1, \ldots, n-1$, namely, the root lattice of type A_n generated by the rows of the $(n-1) \times (n-1)$-Cartan matrix

$$\begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ & & & -1 & 2 \end{bmatrix},$$

where the empty positions represent zeros [5, Planche 1]. In particular, the index of Λ in $\mathbb{Z}^{n-1}$ equals n. For example, if $n = 3$, here is the root lattice Λ (in black) and its two cosets (in gray and white):



Now let Δ_d be the simplex in $\mathbb{R}^n$ with vertices $0, de_1, \ldots, de_n$, for $i = 1, \ldots, n$ let S_i be the set of lattice points in Δ_d that have ith coordinate zero, and set $S_0 := (\mathbb{Z} \times \Lambda) \cap \Delta_d$. Define

$$S := S_1 \cup S_2 \cup \ldots \cup S_n \cup S_0,$$

a subset of the lattice points in Δ_d. We claim that S is connected to 0. Indeed, for each $i = 1, \ldots, n$ the set S_i is connected to 0, and from each point α in S_0 one can walk within S_0 to S_1 by subtracting α_1 times an e_1.

Next, we claim that for each $\alpha \in \Delta_d \cap \mathbb{Z}^n$ there exists a $\beta \in S$ with $\alpha - \beta \in \{0, e_2, \ldots, e_n\}$. Indeed, there is a (unique) β' with this property in $\mathbb{Z} \times \Lambda$. If this β' has nonnegative entries, then set $\beta := \beta' \in S_0$. Otherwise, α itself has a zero entry, say on the ith position, and we set $\beta := \alpha \in S_i$.

Furthermore, for $i = 1, \ldots, n$ the set S_i contains $O(d^{n-1})$ vertices, and S_0 contains $\frac{1}{n} \cdot \frac{1}{n!} d^n + O(d^{n-1})$ vertices. This concludes the construction–note that in the construction of Proposition 4.3 the matrix M_0 has a zero row, so that it represents a compression space with a one-dimensional witness.

For sharpness, assume that $M = M_0 + M_1$ is a uniform determinantal representation of size N such that the singular matrix space represented by M_0 is a compression space with a one-dimensional witness. After a choice of basis of K^n, we may assume that the first row of M_0 is identically zero; write $M_0 = [0|M_0']^T$ accordingly. Let u be the first row of M_1 and write $M_1 = [u|M_1']^T$. Then we have

$$p = \sum_{|\alpha| \leq d} c_\alpha x^\alpha = \det[u|M_0' + M_1'].$$

Let $D_1, \ldots, D_N$ denote the $(N-1) \times (N-1)$ subdeterminants of M_0'. By Lemma 2.6, the space V spanned by these satisfies $F_1 \cdot V \supseteq F_d$. This already gives a lower bound of V equal to $d^n/((n+1)n!) + O(d^{n-1})$. To improve the $n+1$ in the denominator into an n, we observe that by Cramer's rule, the map

$$F_1^N \to K[x_1, \ldots, x_n], \ (\ell_1, \ldots, \ell_N) \mapsto \sum_i (-1)^i \ell_i D_i$$

has every column of M_0' in its kernel. These columns are linearly independent over K (indeed over $K(x_1, \ldots, x_n)$; see Lemma 2.5). We conclude that

$$N \cdot \dim F_1 - (N-1) \geq \dim F_d, \tag{4.1}$$

so that

$$N \geq ((\dim F_d) - 1)/n = d^n/(n \cdot n!) + O(d^{n-1}),$$

as desired. □

In the next section we derive a second general construction of uniform determinantal representations, which we use to prove Theorem 1.3.

5. A second construction. For a while, we believed that the uniform determinantal representations of Theorem 4.5 were optimal. But then we realised that if one relaxes the condition that M_0 represent a compression space with one-dimensional witness to the condition that M_0 represent *some* compression space, smaller-size representations are possible. The basic example is the following.

EXAMPLE 5.1. Let $p = \sum_{i+j \leq 4} c_{ij} x^i y^j$ be the generic polynomial of degree $d = 4$ in $n = 2$ variables. It has the following uniform determinantal representation:

$$p = \det \begin{bmatrix} -x & 1 & & & & & & & \\ & -x & 1 & & & & & & \\ & & -x & 1 & & & & & \\ & & & -x & 1 & & & & \\ c_{00} & c_{10} & c_{20} & c_{30} & c_{40} & -y & & & \\ c_{01} & c_{11} & c_{21} & c_{31} & & 1 & -y & & \\ c_{02} & c_{12} & c_{22} & & & & 1 & -y & \\ c_{03} & c_{13} & & & & & & 1 & -y \\ c_{04} & & & & & & & & 1 \end{bmatrix}, \tag{5.1}$$

where the empty positions denote zeros. Let $M = M_0 + M_1$ be the matrix on the right-hand side. In this case, M_0 represents a compression space with witness $U = \langle e_5, \ldots, e_9 \rangle_K$, which is mapped into $\langle e_6, \ldots, e_9 \rangle_K$.

To verify the identity above without too many calculations, note that the 5 maximal subdeterminants of the 4×5-block with x's are, consecutively, $1, -x, x^2, -x^3, x^4$, and similarly for y. The matrix obtained from M by deleting the column corresponding to x^i and the row corresponding to y^j has determinant $x^i y^j$.

This example extends to a uniform determinantal representation of size $2d+1$ for the generic bivariate polynomial p of degree d. We get $p = \det(M)$, where

$$M = (-1)^d \begin{bmatrix} M_x & 0 \\ L & M_y^T \end{bmatrix},$$

M_x and M_y are $d \times (d+1)$ matrices with 1 on the first upper diagonal and $-x$ and $-y$ respectively on the main diagonal, while L is a $(d+1) \times (d+1)$ triangular matrix such that $\ell_{ij} = p_{j-1,i-1}$ for $i + j \leq d + 2$ and 0 otherwise. Note that we will slightly improve on the size $2d + 1$ in Example 5.3. ♣

Example 5.1 generalises as follows.

PROPOSITION 5.2. *Let $V, W \subseteq K[x_1, \ldots, x_n]$ be subspaces connected to 1 such that $F_1 \cdot V \cdot W \supseteq F_d$. Then there exists a uniform determinantal representation of the generic n-variate polynomial of degree d of size $-1 + \dim V + \dim W$.*

Proof. Set $m_1 := \dim V$ and $m_2 := \dim W$. Consider the matrix

$$M := \begin{bmatrix} M_V & 0 \\ L & M_W^T \end{bmatrix},$$

with M_V and M_W the matrices of sizes $(m_1 - 1) \times m_1$ and $(m_2 - 1) \times m_2$ from Lemma 4.2, and where $L = (\ell_{ij})_{ij}$ is an $m_2 \times m_1$-matrix to be determined. Note that the determinant of M is linear in the entries of L. Indeed, setting $L = 0$ yields the singular matrix M_0, so $\det(M)$ contains no terms of degree 0 in the entries of L. Furthermore, deleting from M two or more of the first m_1 columns from M_V, we end up with a matrix that is singular since, when acting on rows, it maps the span of $\langle e_1, \ldots, e_{m_1-1}\rangle$ into a space of dimension at most $m_1 - 2$, so $\det(M)$ does not contain terms that are of degree > 1 in the entries of L.

Hence the determinant equals $\sum_{ij} \pm \ell_{ij} D_j E_i$ where the D_j are the maximal subdeterminants of M_V and the E_i are the maximal subdeterminants of M_W. By Lemma 4.2 we have $V = \langle D_1, \ldots, D_{m_1}\rangle_K$ and $W = \langle E_1, \ldots, E_{m_2}\rangle_K$. Hence the assumption that $F_1 \cdot V \cdot W \supseteq F_d$ ensures that we can choose the $\ell_{ij} \in F_1$ in such a manner that the determinant of M equals the generic polynomial p. □

Example 5.3. Example 5.1 can be slightly improved to a representation of size $2d - 1$ by taking $V = \langle 1, x, \ldots, x^{d-1}\rangle$ and $W = \langle 1, y, \ldots, y^{d-1}\rangle$; note that, indeed, $F_1 \cdot V \cdot W \supseteq F_d$. A representation of size $2d - 1$ for the polynomial p from (5.1) is

$$p = \det \begin{bmatrix} -x & 1 & & & & & \\ & -x & 1 & & & & \\ & & -x & -1 & & & \\ c_{00} & c_{10} & c_{20} & c_{30} + c_{40}x & -y & & \\ c_{01} & c_{11} & c_{21} + c_{31}x & & 1 & -y & \\ c_{02} + c_{03}y & c_{12} + c_{22}x & & & & 1 & -y \\ c_{13}x + c_{04}y & & & & & & 1 \end{bmatrix}. \tag{5.2}$$

We do not know whether the factor 2 can be improved. ♣

Remark 5.4. A representation of the form (5.2) can also be obtained from the linearisations based on dual basis from [32]. There, linearisations of a univariate polynomial are presented that use the basis of the form $\varphi_i(x)\psi_j(x)$, where φ_i and ψ_j are polynomials. If we use the same approach for a bivariate polynomial with the standard basis $\varphi_i = x^i$ and $\psi_j = y^j$, we get a representation of the form (5.2) up to permutations of rows and columns.

We will now prove our main theorem.

Proof. [Proof of Theorem 1.3, lower bound.] For the lower bound on $N^*(n, d)$, let $M = M_0 + M_1$ be a size-N uniform determinantal representation of the generic n-variate polynomial of degree d. Let D_{ij} be the $(N-1) \times (N-1)$-determinant of the submatrix of M_0 obtained by deleting the entry at position (i, j). Then the image of the linear map

$$\varphi : F_1^{N \times N} \to K[x_1, \ldots, x_n],\ (\ell_{ij})_{i,j} \mapsto \sum_{i,j} (-1)^{i+j} \ell_{ij} D_{ij}$$

contains F_d (see Lemma 2.6).

We claim that φ has a kernel of dimension at least $N(N-1)$. Indeed, fix any row index i_0. If the $D_{i_0,j}$ are all zero, then we obtain an $(n+1)N$-dimensional subspace of $\ker \varphi$ by setting all $\ell_{i,j}$ with $i \neq i_0$ equal to zero and choosing the $\ell_{i_0,j} \in F_1$ arbitrarily. If they are not all zero, then the $N - 1$ rows with indices $i_1 \neq i_0$ are linearly independent over $K(x_1, \ldots, x_n)$ and hence

a fortiori over K. For each such i_1 define $\ell^{(i_1)} \in F_1^{N\times N}$ by

$$\ell_{ij}^{(i_1)} := \begin{cases} (M_0)_{i_1,j} & \text{if } i = i_0, \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

Then, by Cramer's rule, we have $\varphi(\ell^{(i_1)}) = 0$, and these $N-1$ vectors are linearly independent. Hence, for each i_0 we find a subspace of $\ker\varphi$ of dimension at least $N-1$, and these subspaces are linearly independent. Thus we find that

$$N^2(n+1) - N(N-1) = N^2 n + N \geq \dim F_d = \frac{d^n}{n!} + O(d^{n-1}),$$

from which the existence of C_1 follows. □

REMARK 5.5. In the proof of the lower bound we have been a bit more careful than strictly needed: without the discussion of the kernel it follows that $N^2 \geq (\dim F_d)/(n+1)$. But one derives a better constant (for $d \to \infty$) by using the kernel.

For the upper bound, we first give a simple construction for even n. For odd n, a trickier analysis is needed (which also applies in the even case); see below.

Proof. [Proof of Theorem 1.3, upper bound for even n] Assume that $n = 2m$ with $m \in \mathbb{Z}_{\geq 0}$. Let V be the space of polynomials in $x_1, \ldots, x_m$ of degree at most d, and let W be the space of polynomials in $x_{m+1}, \ldots, x_n$ of degree at most d. Then V and W are connected to 1 and we have $F_1 \cdot V \cdot W \supseteq F_d$, so that by Proposition 5.2 we have $N^*(n,d) \leq -1 + \dim V + \dim W$. Now compute

$$\dim V = \dim W = \binom{m+d}{m} = \frac{d^{n/2}}{(n/2)!} + O(d^{m-1}).$$

This implies the existence of C_2 for even n. □

We now give a construction that works for all $n > 2$, for which we thank Aart Blokhuis.

Proof. [Proof of Theorem 1.3, upper bound for $n > 2$] For $i = 0, 1$ let $B_i \subseteq \mathbb{Z}_{\geq 0}$ denote the set of nonnegative integers that can be expressed as $\sum_{j=0}^{e} b_j 2^{2j+i}$ with $b_j \in \{0, 1\}$, i.e., whose binary expansions have ones only at even positions (for $i = 0$, counting the least significant bit as zeroeth position) or only at odd positions (for $i = 1$). Observe that $B_0 + B_1 = \mathbb{Z}_{\geq 0}$ and that both B_0 and B_1 contain roughly $\sqrt{d}$ of the first d nonnegative integers for every d–they have "dimension 1/2". Now set $A_i := B_i^n \subseteq \mathbb{Z}_{\geq 0}^n$ for $i = 0, 1$, so that $A_0 + A_1 = \mathbb{Z}_{\geq 0}^n$ and the number of elements of A_i intersected with a large box $[0, d]^n$ is roughly $d^{n/2}$.

Unfortunately, A_0 and A_1 are not connected to 0. However, we can connect them to 0 as follows. For a lattice point $\alpha \in A_i \setminus \{0\}$ let l be the minimum among the 2-adic valuations of its entries, attained, say, by α_j. Then set $\tilde{\alpha}_j := \alpha_j - 2^l \in B_i$. Setting the remaining coordinates of $\tilde{\alpha}$ equal to those of α we have $\tilde{\alpha} \in A_i$ and

$$\|\alpha - \tilde{\alpha}\|_1 = 2^l.$$

We propose to add to A_i the sequence

$$\alpha - e_j, \alpha - 2e_j, \ldots, \alpha - (2^l - 1)e_j$$

to connect α to $\tilde{\alpha}$. We need to verify, however, that in doing this, A_i retains dimension $n/2$. The fraction of α in (a large box intersected with) A_i for which the minimal valuation is at least l equals roughly $(2^{(-l+i)/2})^n$–after all, the condition is that for each $j = 1, \ldots, n$, α_j has zeros on

the first $(l-i)/2$ positions where it is allowed to have ones. Write $l = i + 2m$. Thus by adding the sequences above, the total increase of A_i is by a factor of at most

$$\sum_{m=0}^{\infty} 2^{2m+i}(2^{-m})^n = \sum_{m=0}^{\infty} 2^{(2-n)m+i}.$$

This is a convergent series as $n > 2$, and hence A_i retains dimension $n/2$.

We have thus constructed subsets $A_0, A_1 \subseteq \mathbb{Z}^n_{\geq 0}$ that satisfy $A_0 + A_1 = \mathbb{Z}^n_{\geq 0}$, are connected to 1, and that contain roughly a constant times $d^{n/2}$ points in each box $[0,d]^n$. For $i = 0, 1$ let V_i be the space spanned by the monomials whose exponent vectors lie in $A_i \cap [0,d]^n$. Then $V_1 \cdot V_2 \supseteq F_d$ and V_1, V_2 are connected to 1, so by Proposition 5.2 there exists a uniform determinantal representation for $p_{n,d}$ of size $\dim V_1 + \dim V_2 = O(d^{n/2})$, as desired. □

Remark 5.6. The construction in the proof is by no means tight. For example, one could also replace α_j by $\alpha_j - 2^l + 2^{l-2} + 2^{l-4} + \ldots + 2^i$, which yields a shorter sequence to be added; and for large l we have also counted additional, shorter sequences since they also have valuation larger than numbers smaller than l. We think that for even n the previous construction, subdividing the variables into two sets of equal size, may lead to a better constant, but we have not verified this.

Example 5.7. Carrying out the construction in the proof for A_0 with $n = 3$, always choosing for j the smallest index of a coordinate of α with minimal valuation, we arrive at the following fractal-like structure (the circles indicate the points of A_0, the black edges show that A_0 is connected to 0):



6. Small n and d. In this section we give several uniform representations of–to our knowledge–the smallest possible size for cases where n and d are small. We start with the two cases where we can compute $N^*(n,d)$ exactly.

Proposition 6.1. $N^*(2,2) = 3$.

Proof. Taking $V = \langle 1, x, y\rangle$ in Proposition 4.3 we see that $N^*(2,2) \leq 3$; this is the representation of Example 1.1. Suppose that a uniform determinantal representation $M = M_0 + M_1$ of size $N = 2$ exists. Then, by Example 3.2, after acting with $\mathrm{SL}_2(K) \times \mathrm{SL}_2(K)$ and transposing if necessary, we may assume that the singular space represented by M_0 is a compression space with a one-dimensional witness. But then (4.1) reads

$$2 \cdot 3 - 1 = N \cdot \dim F_1 - (N-1) \geq \dim F_2 = 6,$$

a contradiction. Hence $N^*(2,2) = 3$. □

Proposition 6.2. $N^*(3,2) = 4$.

Proof. Taking $V = \langle 1, x, y, z\rangle$ in Proposition 4.3 we see that $N^*(3,2) \leq 4$. Suppose that a uniform representation of size $N = 3$ exists. Up to transposition, there are three possibilities for

the singular space $\mathcal{A}$ represented by M_0; see Example 3.2 (where the third compression space is conjugate to the transpose of the first):

1. Assume that $\mathcal{A}$ is a compression space with a one-dimensional witness, so that after acting with $\mathrm{SL}_3(K) \times \mathrm{SL}_3(K)$ we have

$$M_0 = \begin{bmatrix} 0 & 0 & 0 \\ * & * & * \\ * & * & * \end{bmatrix}.$$

Let D_j denote the determinant of the minor of M_0 obtained by deleting the first row and jth column. Then the linear map

$$\Omega : F_1^3 \mapsto K[x, y, z],\ (l_1, l_2, l_3) \mapsto l_1 D_1 - l_2 D_2 + l_3 D_3$$

has $F_2 \subseteq \operatorname{im} \Omega$. Now inequality (4.1) reads

$$3 \cdot 4 - 2 = N \cdot \dim F_1 - (N - 1) \geq \dim F_2 = 10,$$

which holds with equality. This means that, in fact, $\operatorname{im} \Omega$ must *equal* F_2. In particular, D_1, D_2, D_3 must all be of degree one (or else $\operatorname{im} \Omega$ would contain cubic polynomials). The image of Ω depends only on the span $V := \langle D_1, D_2, D_3 \rangle \subseteq F_1$. If $1 \notin V$, then there exists an affine transformation in $\mathrm{AGL}_3(K)$ that maps V into a subspace of $\langle x, y, z \rangle$. Then $1 \notin F_1 \cdot V = \operatorname{im} \Omega$, a contradiction. If $1 \in V$, then after an affine transformation we find $\langle 1 \rangle \subseteq V \subseteq \langle 1, x, y \rangle$. In that case, $z^2 \notin F_1 \cdot V$, another contradiction.

2. Assume that $\mathcal{A}$ is a compression space with a two-dimensional witness, so that after row and column operations we have

$$M_0 = \begin{bmatrix} 0 & 0 & q \\ 0 & 0 & r \\ s & t & * \end{bmatrix},$$

where $q, r, s, t \in F_1$. Write $M_1 = (m_{ij})_{ij}$. Using that $\det(M)$ is assumed to be linear in the c_αs, we find that

$$\det(M) = -m_{11} rt + m_{12} rs + m_{21} qt - m_{22} qs.$$

Consequently, setting $V_1 := \langle q, r \rangle$ and $V_2 := \langle s, t \rangle$, we have $F_1 \cdot V_1 \cdot V_2 \supseteq F_2$. If $1 \notin V_1$, then by acting with a suitable element of $\mathrm{AGL}_3(K)$ we achieve that $V_1 \subseteq \langle x, y, z \rangle$. But then $F_1 \cdot V_1 \cdot V_2 \not\ni 1$. The same applies when $1 \notin V_2$. On the other hand, if $1 \in V_1 \cap V_2$, then by an element in $\mathrm{AGL}_3(K)$ we achieve that $\langle 1 \rangle \subseteq V_1, V_2 \subseteq \langle 1, x, y \rangle$. In that case, $z^2 \notin F_1 \cdot V_1 \cdot V_2$.

3. Finally, assume that $\mathcal{A}$ is conjugate to a space of skew-symmetric matrices, so that after conjugation

$$M_0 = \begin{bmatrix} 0 & q & r \\ -q & 0 & s \\ -r & -s & 0 \end{bmatrix}$$

where $q, r, s \in F_1$. Set $V := \langle q, r, s \rangle \subseteq F_1$. Then the space spanned by the 2×2-determinants of M_0 is $V \cdot V$, of dimension at most 6. Moreover, we have $F_1 \cdot V \cdot V \supseteq F_2$. If $1 \notin V$, then by acting with $\mathrm{AGL}_3(K)$ we achieve that $V \subseteq \langle x, y, z \rangle$, and hence $1 \notin F_1 \cdot V \cdot V$. If, on the other hand, $1 \in V$, then we achieve that $\langle 1 \rangle \subseteq V \subseteq \langle 1, x, y \rangle$, and $z^2 \notin F_1 \cdot V \cdot V$.

In each of these cases we arrive at a contradiction. Consequently, $N^*(3,2) = 4$ as claimed. □

The proofs above use the classification of spaces of small singular matrices in an essential manner, as well as the action of $\mathrm{AGL}_n(K)$ on uniform determinantal representations. We conjecture that $N^*(4,2) = 5$, and that this can still be proved in the same manner, using the classification of 4×4-singular matrix spaces from [13]. But as Theorem 3.4 shows, fundamentally new ideas will be needed to prove lower bounds in larger situations.

For some pairs of small n and d we now give the smallest uniform representations that we have been able to find. For the constructions we use Proposition 5.2 with subspaces $V, W \subseteq K[x_1, \ldots, x_n]$ spanned by the monomials and connected to 1. First, we give in Table 6.1 the minimal sizes known to us of uniform determinantal representations for some small values of n and d.

TABLE 6.1

Minimal known sizes of uniform determinantal representations we have been able to construct for n-variate polynomials of degree d; cf. Table 6.2.

n	$d=2$	$d=3$	$d=4$	$d=5$	$d=6$	$d=7$	$d=8$	$d=9$
2	3	5	7	9	11	13	15	17
3	4	7	10	14	18	22	27	34
4	5	9	14	19	26	34	44	
5	6	11	18	26				
6	7	13	22	33				
7	8	15	27	39				
8	9	17	32					

The corresponding representations for the entries in Table 6.1 for $n = 2$, which are of size $2d-1$, are given in Example 5.3. For $d = 2$ we take $V = \langle 1, x_1, \ldots, x_n\rangle$ and $W = \langle 1\rangle$, therefore $N^*(n,2) \le n+1$, while for $d = 3$ we can take $V = W = \langle 1, x_1, \ldots, x_n\rangle$, and hence $N^*(n,3) \le 2n-1$. In Table 6.2 we give sets V and W for the remaining nonzero entries in Table 6.1. The subspaces V and W have the form $V = V_0 \cup V_1$ and $W = W_0 \cup W_1$, where

$$\begin{aligned} V_0 &= \langle 1, x_1, \ldots, x_n, \ldots, x_1^e, \ldots, x_n^e\rangle, \\ W_0 &= \langle 1, x_1, \ldots, x_n, \ldots, x_1^f, \ldots, x_n^f\rangle \end{aligned} \tag{6.1}$$

for $e = \lceil (d-1)/2\rceil$ and $f = \lfloor (d-1)/2\rfloor$, which yields $d-1 = e+f$. For clarity and brevity, the variables x, y, z, w, u, v, q, s in Table 6.2 stand for $x_1, \ldots, x_8$, respectively.

EXAMPLE 6.3. To show how things get complicated, let us consider the construction for $d = 4$. We take $V = \langle 1, x_1, \ldots, x_n, x_1^2, \ldots, x_n^2\rangle$ and

$$W = \langle 1, x_1, \ldots, x_n, x_{\alpha_1}x_{\beta_1}, \ldots, x_{\alpha_m}x_{\beta_m}\rangle,$$

where $1 \le \alpha_i < \beta_i \le n$ and m is as small as possible. If we take all possible pairs $x_\alpha x_\beta$, then clearly $F_1 \cdot V \cdot W \supseteq F_4$, while on the other hand, when $m = 0$, $F_1 \cdot V \cdot W$ does not contain any monomials of the form

$$x_i\, x_j\, x_k\, x_\ell \tag{6.2}$$

for $1 \le i < j < k < \ell \le n$. We need a minimal set of $x_\alpha x_\beta$ to cover all possible monomials (6.2), which is related to the following covering problem.

Given positive integers $r \le k \le n$, we say that a system S of r-subsets (called blocks) of $\{1, \ldots, n\}$ is called a Turán (n,k,r)-system if every k-subset of $\{1, \ldots, n\}$ contains at least one block from S [34]. The minimum size of S is called the Turán number $T(n,k,r)$.

TABLE 6.2

List of monomials in V_1 and W_1 that, together with V_0 and W_0 from (6.1), lead to uniform representations of sizes as in Table 6.1.

n	d	V_1	W_1
3	4	–	–
3	5	–	xy
3	6	–	xy, x^2y
3	7	–	x^2y, y^2z, z^2x
3	8	–	x^2y, y^2z, z^2x, x^2y^2, z^2w^2
3	9	x^3y, y^3z, z^3x	x^2y, x^2z, y^2z, x^2y^2, x^2z^2, y^2z^2
4	4	–	xy
4	5	–	xy, zw
4	6	x^2y, y^2z, z^2w	xy, x^2y, zw
4	7	x^2y, y^2x, z^2w, w^2x, xy	x^2z, xz^2, y^2w, yw^2
4	8	x^2y, x^2y^2, z^2x, x^3y, y^3z, z^3w, w^3x	xy, xyz, xyw, y^2z, z^2w, w^2x, w^2y, x^2z
5	4	–	xy, zw
5	5	xy, yz, zw	wu, xu
6	4	–	xy, zw, uv
6	5	xy, zw, uv, wy	yz, wu, xv, xz
7	4	–	xy, zw, uv, xq, yq
7	5	xy, zw, uv, wy, qu	yz, wu, vq, xz, wx
8	4	–	xy, yz, xz, wu, wv, uv, qs

In our case, additional terms $x_{\alpha_1}x_{\beta_1}, \dots, x_{\alpha_m}x_{\beta_m}$ form a Turán $(n, 4, 2)$-system. While for most cases only upper and lower bounds for $T(n, k, r)$ are known, Turán proved that

$$T(n,4,2) = mn - 3\,\frac{m(m+1)}{2}, \tag{6.3}$$

where $m = \lfloor n/3 \rfloor$. To obtain the minimal set one has to divide $\{1, \dots, n\}$ into three nearly equal groups (their sizes do not differ for more than one) and then take all pairs $x_\alpha x_\beta$ such that α and β belong to the same group. As a result, such construction gives a uniform representation of size N, where $N = \frac{1}{6}\,n^2 + O(n)$, which therefore implies $N^*(n, 4) \le \frac{1}{6}\,n^2 + O(n)$. ♣

7. Matrix polynomials. Suppose that we have a uniform representation M of $p_{n,d}$ as in (2.1), and write

$$M = M_0 + M_1 = M_0 + \sum_{|\alpha| \le d} c_\alpha M_\alpha, \tag{7.1}$$

where each $M_\alpha \in F_1^{N \times N}$. Now consider the matrix polynomial (cf. (2.1))

$$P_{n,d} = \sum_{|\alpha| \le d} x^\alpha C_\alpha,$$

where C_α is a $k \times k$ matrix. We will show that under certain assumptions we can construct from M a matrix $\widetilde{M}$ that represents $P_{n,d}$ in the sense that $\det(\widetilde{M}) = \det(P_{n,d})$. We obtain $\widetilde{M}$ from M in the following way. Each element of the form $\alpha + \beta x + \gamma y$ is replaced by the $k \times k$ matrix $(\alpha + \beta x + \gamma y)I_k$, where I_k is the $k \times k$ identity, and each c_α is replaced by the matrix C_α.

THEOREM 7.1. *Let (7.1) be a uniform representation of the generic polynomial (2.1) of degree d in n variables and assume that there exist matrices Q and Z, whose elements are*

polynomials in $x_1, \ldots, x_n$, such that $\det(Q) = \det(Z) = 1$, *and QMZ is a triangular matrix with one diagonal element equal to $p_{n,d}$ and all other diagonal elements equal to 1. Then*

$$\widetilde{M} = M_0 \otimes I_k + \sum_{|\alpha| \le d} M_\alpha \otimes C_\alpha$$

is a representation for the matrix polynomial $P_{n,d}$, i.e., $\det(\widetilde{M}) = \det(P_{n,d})$.

Proof. It is easy to see that $(Q \otimes I_k)\widetilde{M}(Z \otimes I_k)$ is a block triangular matrix with one diagonal block $P_{n,d}$ while all other diagonal blocks are equal to I_k. Since $\det(Q \otimes I_k) = \det(Z \otimes I_k) = 1$, it follows that $\det(\widetilde{M}) = \det(P_{n,d})$. □

Example 7.2. Theorem 7.1 applies to the uniform representation (5.1). Indeed, take

$$Q = \begin{bmatrix} 1 & & & & & & & & \\ & \ddots & & & & & & & \\ & & 1 & & & & & & \\ & & & 1 & y & y^2 & y^3 & y^4 \\ & & & & 1 & y & y^2 & y^3 \\ & & & & & 1 & y & y^2 \\ & & & & & & 1 & y \\ & & & & & & & 1 \end{bmatrix}, \qquad Z = \begin{bmatrix} 1 & & & & & & & \\ x & 1 & & & & & & \\ x^2 & x & 1 & & & & & \\ x^3 & x^2 & x & 1 & & & & \\ x^4 & x^3 & x^2 & x & 1 & & & \\ & & & & & 1 & & \\ & & & & & & \ddots & \\ & & & & & & & 1 \end{bmatrix},$$

then

$$QMZ = \begin{bmatrix} & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ p & \times & \times & \times & c_{40} & & & \\ \times & \times & \times & c_{31} & & 1 & & \\ \times & \times & c_{22} & & & & 1 & \\ \times & c_{13} & & & & & & 1 \\ c_{04} & & & & & & & \end{bmatrix}.$$

It is easy to see that there exist permutation matrices P_L and P_R such that

$$P_L(QMZ)P_R = \begin{bmatrix} 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ c_{40} & \times & \times & \times & p & & & \\ & \times & \times & c_{31} & \times & 1 & & \\ & \times & c_{22} & & \times & & 1 & \\ & c_{13} & & & \times & & & 1 \\ & & & & c_{04} & & & \end{bmatrix}$$

is triangular and has the diagonal which satisfies Theorem 7.1. Therefore, we can apply (5.1) for matrix polynomials by using block matrices. This can be generalised to a uniform representation of size $2d + 1$ of the form (5.1). In a similar way we can show that this also holds for representations of the form (5.2) of size $2d - 1$. ♣

Unfortunately, not all uniform determinantal representations induce a determinantal representation of a general matrix polynomial in this manner. As a counterexample, let M be such a uniform determinantal representation of the polynomial $p_{n,d}$, $|\alpha|, |\beta| \leq d$, and construct a representation of larger size

$$M' = \begin{bmatrix} M & 0 \\ 0 & N \end{bmatrix}, \qquad \text{with} \quad N = \begin{bmatrix} 0 & c_\alpha & c_\beta & 1 \\ -c_\alpha & 0 & 1 & 0 \\ -c_\beta & -1 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{bmatrix}.$$

Then $\det(M') = \det(M)\det(N) = p_{n,d}(1 + c_\alpha c_\beta - c_\beta c_\alpha) = p_{n,d}$, but $\widetilde{M'}$ is not a representation for the matrix polynomial $P_{n,d}$ as the coefficient matrices C_α and C_β do not commute in general. This motivates the following definition.

DEFINITION 7.3. *A uniform determinantal representation M is* minimal *if there do not exist constant matrices P and Z such that* $\det(P) = \det(Z) = 1$ *and*

$$QMZ = \begin{bmatrix} * & * \\ 0 & M_2 \end{bmatrix}, \qquad \textit{where } M_2 \textit{ is square with} \quad \det(M_2) = 1.$$

We speculate that each minimal uniform representation gives rise to a representation for a matrix polynomial.

8. Numerical experiments. Recently, a new numerical approach for computing roots of systems of bivariate polynomials was proposed in [31]. The main idea is to treat the system as a two-parameter eigenvalue problem using determinantal representations.

Suppose that we are looking for roots of a system of bivariate polynomials

$$\begin{aligned} p &= \textstyle\sum_{i+j\leq d_1} \alpha_{ij} x^i y^j = 0, \\ q &= \textstyle\sum_{i+j\leq d_2} \beta_{ij} x^i y^j = 0, \end{aligned} \tag{8.1}$$

where p and q are polynomials of degree d_1 and d_2 over $\mathbb{C}$. Let $P = A_0 + xA_1 + yA_2$ and $Q = B_0 + xB_1 + yB_2$, where $A_0, A_1, A_2 \in \mathbb{C}^{N_1 \times N_1}$ and $B_0, B_1, B_2 \in \mathbb{C}^{N_2 \times N_2}$, with $\det(P) = p$ and $\det(Q) = q$, be determinantal representations of p and q, respectively. Then a root (x, y) of (8.1) is an eigenvalue of the two-parameter eigenvalue problem

$$\begin{aligned} (A_0 + xA_1 + yA_2)\, u &= 0, \\ (B_0 + xB_1 + yB_2)\, v &= 0, \end{aligned} \tag{8.2}$$

where $u \in \mathbb{C}^{N_1}$ and $v \in \mathbb{C}^{N_2}$ are nonzero vectors. The standard way to solve (8.2) is to consider a joint pair of generalized eigenvalue problems [1]

$$\begin{aligned} (\Delta_1 - x\Delta_0)\, w &= 0, \\ (\Delta_2 - y\Delta_0)\, w &= 0, \end{aligned} \tag{8.3}$$

where

$$\Delta_0 = A_1 \otimes B_2 - A_2 \otimes B_1, \quad \Delta_1 = A_2 \otimes B_0 - A_0 \otimes B_2, \quad \Delta_2 = A_0 \otimes B_1 - A_1 \otimes B_0,$$

and $w = u \otimes v$.

In this particular application we can expect that the pencils in (8.3) are singular, i.e., $\det(\Delta_1 - x\Delta_0) \equiv 0$ and $\det(\Delta_1 - y\Delta_0) \equiv 0$. Namely, by Bézout's theorem a generic system (8.1) has $d_1 d_2$ solutions, while a generic problem (8.2) has $N_1 N_2$ eigenvalues. Unless

$(d_1, d_2) = (N_1, N_2)$, both pencils in (8.3) are singular. In this case we first have to apply the staircase algorithm from [26] to extract the finite regular eigenvalues. The method returns smaller matrices $\widetilde{\Delta}_0$, $\widetilde{\Delta}_1$, and $\widetilde{\Delta}_2$ (of size $d_1 d_2 \times d_1 d_2$ for a generic (8.1)) such that $\widetilde{\Delta}_0$ is non-singular and $\widetilde{\Delta}_0^{-1}\widetilde{\Delta}_1$ and $\widetilde{\Delta}_0^{-1}\widetilde{\Delta}_2$ commute. From

$$(\widetilde{\Delta}_1 - x\widetilde{\Delta}_0)\,\widetilde{w} = 0,$$
$$(\widetilde{\Delta}_2 - y\widetilde{\Delta}_0)\,\widetilde{w} = 0,$$

we compute the eigenvalues (x, y) using a variant of the QZ algorithm [17] and thus obtain the roots of (8.1).

The above approach is implemented in the Matlab package BiRoots [30] together with the two determinantal representations from [31]. The first one, to which we refer as `Lin1`, is a uniform one from Example 4.4 of size $\frac{1}{4}\,d^2 + O(d)$ for a polynomial of degree d. The second one, which we refer to as `Lin2`, is not uniform and involves some computation to obtain a smaller size $\frac{1}{6}\,d^2 + O(d)$. Although the construction of `Lin2` is more time consuming, this pays off later, when the staircase algorithm is applied to (8.3).

Table 8.1 shows the sizes of determinantal representations for polynomials of small degree. As expected, the new uniform determinantal representation of size $2d - 1$, to which we refer as `MinUnif`, returns smaller matrices, which reflects later in faster computational times. It is also important that `Lin1` and `MinUnif` return real matrices for polynomials with real coefficients, which is not true for `Lin2`.

TABLE 8.1
Size of the matrices for `Lin1` *and* `Lin2` *for bivariate polynomials ($n = 2$) and various degrees d.*

Method	$d=3$	$d=4$	$d=5$	$d=6$	$d=7$	$d=8$	$d=9$	$d=10$	$d=11$	$d=12$
`Lin1`	5	8	11	15	19	24	29	35	41	48
`Lin2`	3	5	8	10	13	17	20	24	29	34
`MinUnif`	5	7	9	11	13	15	17	19	21	23

It was reported in [31] that the determinantal representation approach for solving systems of bivariate polynomials is competitive for polynomials of degree 9 or less. As we show below, the new uniform representation `MinUnif` extends this to degree 15 and, in addition, performs better than the existing representations for polynomials of degree 6 or more.

In [31] the approach was compared numerically to the following state-of-the art numerical methods for polynomial systems: `NSolve` in Mathematica 9 [41], BertiniLab 1.4 [29] running Bertini 1.5 [3], NAClab 3.0 [42], and `PHCLab` 1.04 [15] running PHCpack 2.3.84, which turned out as the fastest of these methods. To show the improved performance of the new determinantal representation, we compare `MinUnif` to `Lin1`, `Lin2`, and `PHCLab` in Table 8.2. For each d we run the methods on the same set of 50 real and 50 complex random polynomial systems of degree d and measure the average time. For `Lin1` and `MinUnif`, where determinantal representations have real matrices for real polynomials, we report separate results for polynomials with real and complex coefficients. The timings for `Lin1` and `Lin2` are given only for $n \leq 10$ as for larger n these two linearisations are no longer competitive.

Of course, the computational time is not the only important factor, we also have to consider the accuracy and reliability. In each step of the staircase algorithm a rank of a matrix has to be estimated numerically, which is a delicate task. After several steps it may happen that the gap between the important singular values and the meaningful ones that should be zero in exact computation, virtually disappears. In such case the algorithm fails and does not return any roots. As the number of steps in the staircase algorithm increases with degree of the polynomials, such

TABLE 8.2
Average computational times in milliseconds for `Lin1`*,* `Lin2`*,* `MinUnif`*, and* `PHCLab` *for random full bivariate polynomial systems of degree* 3 *to* 15*. For* `Lin1` *and* `MinUnif` *separate results are included for real* ($\mathbb{R}$) *and complex polynomials* ($\mathbb{C}$)*.*

d	`Lin1` ($\mathbb{R}$)	`Lin1` ($\mathbb{C}$)	`Lin2`	`PHCLab`	`MinUnif` ($\mathbb{R}$)	`MinUnif` ($\mathbb{C}$)
3	6	8	4	116	6	7
4	9	11	6	130	12	13
5	20	26	13	151	18	20
6	39	71	28	174	27	27
7	96	160	51	217	36	44
8	205	395	118	264	59	74
9	467	1124	279	329	95	125
10	1424	3412	600	414	147	221
11				538	248	354
12				650	361	530
13				911	592	740
14				1142	842	1148
15				1531	1237	1835

problems occur more often for polynomials of large degree. A heuristic that usually helps in such cases is to apply the algorithm on a transformed system

$$\begin{aligned}\widetilde{p} &:= \quad cp + sq = 0,\\ \widetilde{q} &:= -sp + cq = 0\end{aligned}$$

for random c and s such that $c^2+s^2 = 1$. As this transformation does not change the conditioning of the roots, we can conclude that the difficulties with the staircase algorithm are not directly related to the conditioning. The trick does not work every time, and it seems that for some systems the only way to make the determinantal representation approach to work is to increase the machine precision.

We can apply the same approach to systems of polynomials in more than two variables. However, since the size of the corresponding Δ matrices is the product of sizes of all representations, this is competitive only for $n = 3$ and $d \leq 3$. For a comparison, if we have a system of three polynomials in three variables of degree 3, then the size of the Δ matrices is 343×343. For degree 4 the size increases to 1000×1000 and PHCpack is faster. Finally, for $n = 4$ and the smallest nontrivial $d = 2$ we already get Δ matrices of size 625×625 and the method is not efficient.

9. Outlook. We have introduced *uniform determinantal representations*, which rather than representing a single polynomial as the determinant of a matrix of affine-linear forms, represent all polynomials of degree at most d in n variables as such a determinant. We have seen that in the bivariate case, these determinantal representations are useful for numerically solving bivariate systems of equations; and in the general multivariate case we have determined, up to constants, the asymptotic behaviour of $N^*(n, d)$, the minimal size of such a representation, for n fixed and $d \to \infty$.

We now summarise several results that have been shown in the paper.

- For fixed n and $d \to \infty$, $N^*(n, d) \sim d^{n/2}$, see Theorem 1.3. This is a noticeable improvement on [33], where an asymptotic rate of $N^*(n, d) \sim d^n$ is shown, with the remark that the representation in [33] are symmetric. However, symmetry currently cannot be exploited by methods that compute roots of multivariate polynomial systems.
- For fixed odd d and $n \to \infty$, $N^*(n, d) \sim n^{(d-1)/2}$, which is the same rate as Quarez [33], who also manages to get symmetric representations. For fixed even d and $n \to \infty$,

$N^*(n,d) \lesssim n^{d/2}$, which again is the same rate as in [33]. However, we have a slightly smaller lower bound for the asymptotic rate of $n^{(d-1)/2}$.

- Tables 6.1 and 6.2 give constructions for the smallest representations that we have been able to find for some small values of n and d.
- $N^*(n,2) \le n+1$; cf. Table 6.1.
- $N^*(n,3) \le 2n+1$; cf. Table 6.1.
- $N^*(n,4) \le \frac{1}{6}n^2 + O(n)$; see Example 6.3.
- $N^*(2,d) \le 2d-1$; cf. Table 6.1. Note that this result satisfies Dixon [10] up to an asymptotic factor 2 whereby no computations are necessary for the determinantal representation. In particular, it is a major improvement on the $\sim \frac{1}{4}d^2$ of [33, 31].
- Due to the smaller sizes of the representations, the numerical approach for bivariate polynomials ($n=2$) is competitive to (say) Mathematica for degree d up to $d \approx 15$ (see Section 8); this in contrast to $d \approx 9$ as obtained in [31].
- Under some conditions, the results carry over to the case of matrix coefficients (see Section 7).

There are still many interesting open questions, both of intrinsic mathematical interest and of relevance to polynomial system solving. First of all, in a situation where the degree d is fixed and the number n of variables grows, what is the asymptotic behaviour of $N^*(n,d)$? Although we expect that the above inequalities are equalities, whether this holds is still open. For general fixed d, the proof of Theorem 1.3 yields a lower bound which is a constant (depending on d) times $n^{(d-1)/2}$. For odd d we obtain a matching upper bound (with a different constant) by using Proposition 5.2 with $V = W = F_{(d-1)/2}$. However, for even d we only know how to obtain $O(n^{d/2})$. We remark that this latter is (up to a constant) the same bound as obtained in [33, Thm. 4.4] for *symmetric* uniform representations.

Second, in the case of fixed n and varying d studied in this paper, what are the best constants in Theorem 1.3? More specifically, for fixed n, does $\lim_{d\to\infty} \frac{N^*(n,d)}{d^{n/2}}$ exist, and if so, what is its value?

Third, how can our techniques for upper bounds and lower bounds be further sharpened? Can singular matrix spaces *other* than compression spaces be used to obtain tighter upper bounds (constructions) on $N^*(n,d)$? Can the action of the affine group be used more systematically to find lower bounds on $N^*(n,d)$?

Fourth, is it true that each minimal uniform representation gives rise to a representation of the corresponding matrix polynomial (cf. Section 7)?

Finally, we have restricted our attention to matrices that, apart from being affine-linear in $x_1,\dots,x_n$, are also affine-linear in the coefficients c_α. Our proofs give the same asymptotic behaviour (with different constants) if we require, in addition, that no quadratic terms $c_\alpha x_i$ may occur in M. If, instead, we relax the condition that M be affine-linear in the c_α to a *polynomial* dependence on the c_α, then the same bounds still apply; see Remark 2.7. But what if we relax this to *rational* dependence of M on the c_α? Given that $p_{n,d}$ is only linear in the c_α it seems unlikely that allowing M to be rational in the c_α we would gain anything, but we currently do not know how to formalise this intuition. On the other hand, in cases where a (non-uniform) determinantal representation of size d is known to exist for every (or sufficiently general) polynomials of degree d in d variables (e.g., in the case of plane curves), it follows that this representation can be chosen to have entries *algebraic* in the c_α. This observation rules out approaches aimed at proving lower bounds in a too general setting.

REFERENCES

[1] F. V. Atkinson, *Multiparameter Eigenvalue Problems*, Academic Press, New York (1972).
[2] A. Beauville, Determinantal hypersurfaces, Mich. Math. J. 48 (2000), 39–64.
[3] D. J. Bates, J. H. Hauenstein, A. J. Sommese, and C. W. Wampler, Bertini: Software for Numerical Algebraic Geometry, available at `bertini.nd.edu`.
[4] A. Boralevi, D. Faenzi and E. Mezzetti, Linear spaces of matrices of constant rank and instanton bundles, Adv. Math. 248 (2013), 895–920.
[5] N. Bourbaki, *Groupes et Algèbres de Lie, Chapitres IV, V et VI*, Éléments de Mathématique XXXIV, Hermann, Paris (1968).
[6] P. Bürgisser, C. Ikenmeyer, and G. Panova, No occurrence obstructions in geometric complexity theory, `arXiv:1604.06431` (2016).
[7] P. Brändén, Obstructions to determinantal representability, Adv. Math. 226(2), (2011), 1202–1212.
[8] L. E. Dickson, Determination of all general homogeneous polynomials expressible as determinants with linear elements, Trans. Amer. Math. Soc. 22 (1921), 167–179.
[9] J. Dieudonné, Sur une généralisation du groupe orthogonal à quatre variables, Arch. Math., Oberwolfach (1949), 1:282287.
[10] A. Dixon, Note on the reduction of a ternary quartic to a symmetrical determinant, Proc. Camb. Phil. Soc. 11 (1902), 350–351.
[11] J. Draisma, Small maximal spaces of non-invertible matrices, Bull. Lond. Math. Soc. 38(5) (2006), 764–776.
[12] D. Eisenbud and J. Harris, Vector spaces of matrices of low rank, Adv. Math. 70(2) (1988), 135–155.
[13] P. Fillmore, C. Laurie, and H. Radjavi, On matrix spaces with zero determinant, Lin. Multilin. Algebra, 18 (1985), 255–266.
[14] H. Flanders, On spaces of linear transformations with bounded rank, J. Lond. Math. Soc. 37 (1962), 10–16.
[15] Y. Guan and J. Verschelde, PHClab: A MATLAB/Octave interface to PHCpack. In: M. Stillman, J. Verschelde, and N. Takayama (eds), *Software for Algebraic Geometry*, volume 148 of The IMA Volumes in Mathematics and its Applications, Springer, New York (2008), 15–32.
[16] J. W. Helton and V. Vinnikov, Linear matrix inequality representation of sets, Commun. Pure Appl. Math. 60(5) (2007), 654–674.
[17] M. E. Hochstenbach, T. Košir, and B. Plestenjak, A Jacobi–Davidson type method for the nonsingular two-parameter eigenvalue problem, SIAM J. Matrix Anal. Appl. 26 (2005), 477–497.
[18] J. Hüttenhain and P. Lairez, The boundary of the orbit of the 3 by 3 determinant polynomial, `arXiv:1512.02437` (2015).
[19] B. Ilic and J. M. Landsberg, On symmetric degeneracy loci, spaces of symmetric matrices of constant rank and dual varieties, Math. Ann. 314(1) (1999), 159–174.
[20] Y. Ishitsuka and T. Ito, On the symmetric determinantal representations of the Fermat curves of prime degree, Int. J. Number Theory 12(4) (2016), 955–967.
[21] J-B. Lasserre, M. Laurent, B. Mourrain, Ph. Rostalski, and Ph. Trébuchet, Moment matrices, border bases and real radical computation, J. Symb. Comput. 51 (2013), 63–85.
[22] A. S. Lewis, P. A. Parrilo, and M. V. Ramana, The Lax conjecture is true, Proc. Am. Math. Soc. 133(9), (2005), 2495–2499.
[23] L. Manivel and E. Mezzetti, On linear spaces of skew-symmetric matrices of constant rank, Manuscripta Math. 117(3) (2005), 319–331.
[24] The MathWorks, Inc., Matlab, Natick, Massachusetts, United States.
[25] A. Muhič and B. Plestenjak, On the singular two-parameter eigenvalue problem, Electron. J. Linear Algebra 18 (2009), 420–437.
[26] A. Muhič and B. Plestenjak, On the quadratic two-parameter eigenvalue problem and its linearization, Linear Algebra Appl. 432 (2010), 2529–2542.
[27] K. D. Mulmuley and M. Sohoni, Geometric complexity theory. I: An approach to the P vs. NP and related problems, SIAM J. Comput. 31(2) (2001), 496–526.
[28] K. D. Mulmuley and M. Sohoni, Geometric complexity theory II: Towards explicit obstructions for embeddings among class varieties, SIAM J. Comput. 38(3) (2008), 1175–1206.
[29] A. Newell, BertiniLab: toolbox for solving polynomial systems, MATLAB Central File Exchange, `www.mathworks.com/matlabcentral/fileexchange/48536-bertinilab`.
[30] B. Plestenjak, BiRoots, MATLAB Central File Exchange, `www.mathworks.com/matlabcentral/fileexchange/54159-biroots` (2015).
[31] B. Plestenjak and M. E. Hochstenbach, Roots of bivariate polynomial systems via determinantal representations, SIAM J. Sci. Comput. 38 (2016), A765–A788.

[32] L. Robol, R. Vandebril, and P. van Dooren, A framework for structured linearizations of matrix polynomials in various bases, arXiv:1603.05773 (2016).
[33] R. Quarez, Symmetric determinantal representation of polynomials, Linear Algebra Appl. 436(9) (2012), 3642–3660.
[34] A. Sidorenko, What we know and what we do not know about Turán numbers, Graphs Combin. 11 (1995), 179–199.
[35] C. de Seguins Pazzis, Large affine spaces of matrices with rank bounded below, Linear Algebra Appl. 437(2) (2012), 499–518.
[36] J. Sylvester, On the dimension of spaces of linear transformations satisfying rank conditions, Linear Algebra Appl. 78 (1986), 1–10.
[37] L. G. Valiant, The complexity of computing the permanent, Theoret. Comput. Sci. 8(2) (1979), 189–201.
[38] J. Verschelde, Algorithm 795: PHCpack: a general-purpose solver for polynomial systems by homotopy continuation, ACM Trans. Math. Softw., 25 (1999), 251–276.
[39] D. G. Wagner, Multivariate stable polynomials: theory and applications, Bull. Am. Math. Soc., New Ser. 48(1), (2011), 53–84.
[40] R. Westwick, Spaces of matrices of fixed rank, Lin. Multilin. Algebra 20 (1987), 171–174.
[41] Wolfram Research, Inc., Mathematica, Version 9.0, Champaign, Illinois (2012).
[42] Z. Zeng and T.-Y. Li, NAClab: a Matlab toolbox for numerical algebraic computation, ACM Commun. Comput. Algebra 47 (2013), 170–173.